

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

## Northern Blot: 7 Bacterial Species Probed with Conserved 22mer from 4.5S

---



Hybridized with Ad4.5S13Vnf:

- RT
- 120V
- 5% gel
- 8M urea

**Fig. 1**

## Northern Blot to compare abundance of 5S and 4.5S





Fig. 2

## Detection of 4.5S RNA in Cultured E.coli by Adapter-Capture and Fluorescent Sandwich Probe

---



3λ /20 RNA from 1 ml of overnight culture

**Fig. 3**

## In-Gel Strand Displacement




Fig. 4

## Adapters for analysis of RNA on Universal Capture Gels.





Fig. 5



**Fig. 6**




Fig. 7